

Disposal of Trees Affected by the Pine Beetle:

The Dilemma and why Air Curtain Burners Should Be Used

By N. Fuhrmann

The Problem

Forest vegetation management in many areas throughout North America is confronted with the dilemma of how to prevent the destruction of trees by pests, such as beetles, preserve threatened trees and safely remove and destroy unsalvageable and dead trees killed by the pest. The latter problem is addressed here and it is determined that the best disposal option for both freshly felled and dead beetle infested trees is by means of above-ground air curtain burners.

The primary objective is two-fold: (1) freshly felled beetle infested trees must be disposed of in such a manner as not to foster cross-contamination, the spreading of the disease problem to healthy trees and (2) to remove dead trees under forest fuels management initiatives, as the dead standing, leaning and fallen trees otherwise would pose a wildfire hazard, most importantly near or in the wildland-urban interface where well managed defensible zones are imperative to protect residential dwellings, businesses and other structures.

The secondary objective is to devise a method that will (a) economically and efficiently accomplish the goals set forth above and (b) in an environmentally friendly and safe way.

Background

Trees can be stricken with a number of diseases, including pest or fungal inflictions that may adversely affect their rate of growth, strength, longevity, reproduction, impact on the tree, flora and fauna community in which they stand and also their suitability for commercial harvesting. In recent years North American forests, parks and private woodlots have suffered immensurable losses from trees that were destroyed by beetle pests, worms or fungi. For example, oak trees and rhododendron have been destroyed by *Sudden Oak Death Syndrome (SOD)* in California, Oregon and other areas. A variety of beetles, some indigenous to sections of North America, i.e. Colorado, others brought into the North American Continent from foreign countries usually by sloppy ocean vessel cargo management practices, such as the Asian Longhorned Beetle, have killed millions of trees, often causing eradication of patches of forests multiple square miles in size.

Society has a high interest and responsibility in managing and preserving the world's forests for many reasons, above all, because trees are vital for the very existence of mankind considering the balance that photosynthesis provides to the air we breathe. Healthy forests are a "carbon sink." Unfortunately, recent studies have shown that the magnitude of decaying vegetative waste resulting from beetle kills have actually reversed some forest areas from being a carbon sink to a contributor to greenhouse gas emissions¹.

Forests provide a habitat for other vegetation and animals of a myriad of species. Forests made it possible for mankind to evolve as it has from providing shelter, building materials and firewood to hunting grounds. Finally, one cannot overlook the beauty of our trees and the forests that they comprise. Their recreational value should not be underestimated.

Non-government and non-profit endeavors that are people-driven, such as the *Champion Tree Project*² supported by the USDA, are of paramount significance, as they raise public awareness regarding the serious issues that threaten our forests and trees and they provide educational platforms to help prepare young people to better understand the steps required to safeguard our forests for future generations and to cope with the immense dangers facing our environment in general.

About Beetles in the Forest

It should be emphasized that generally most beetles in the forest are not harmful, but rather are a vital part of the forest's ecosystem. It is often thought that the trees in our country are attacked mainly by invasive species, pests that have been imported from foreign lands and that do not have natural local predators. However, there are native beetle species as well that are responsible for massive tree mortalities.

In Colorado, for example, the *mountain pine beetle* (MPB) that is causing extensive damage at the present time and has destroyed countless numbers of trees in recent years, is native to the forests of Western North America.

It is the abundance and concentration of the beetle population that result in the massive destruction of trees. The main reason for providing the basis for such beetle abundance is the fact that many forests in the US and Canada are severely overgrown and, as a result, are weakening their trees, because natural low-severity forest fires have not cleared out the excessive abnormal underbrush and frail trees for decades or even a century. Ever since the US Government implemented wildfire suppression mandates more than 85 years ago, the natural process of clearing out ladder fuels (underbrush or

understory) and dead trees has been disturbed and forest fires have thus become much more severe³. For millions of years, naturally occurring wildfires have not only created a balance between various species of healthy trees, fragile trees and pests, but also provided a biological necessity for certain trees. For example, the *Lodgepole Pine* depends on heat from wildfires to open its cones to release the seeds⁴, and the *California Redwood* also needs fire to survive, as without it redwoods will not properly repopulate⁵.

Trees that have been injured or otherwise weakened from droughts, unnatural high-severity wildfires, etc. are most vulnerable for beetle infestations; however, otherwise healthy appearing trees are also succumbing to an overly large beetle population, to some extent due to the adverse effect of the abnormally dense understory forest growth which makes the stands more susceptible to pest attacks.

It must furthermore be noted that beetles are selective in their host trees. Not all "bark beetles" invade all species of "pine trees"; they typically choose specific tree species as hosts. This is important to understand when devising eradication plans, as one type of pine may be sickened by a particular beetle attack whereas another species of pine tree next to it is just fine and must be left alone^{6;7}.

Eradication of (Harmful) Beetles

Beetle eradication methods are rather limited in scope: chemical pesticides that are aimed at killing the beetle and the underlying larvae are used to salvage healthy trees in danger of beetle infliction. This approach, albeit costly, makes sense in parks, residential areas and private woodlots where selective treatment is realistic, but not in vast forest lands.

The only alternative is to stem the spread of the beetle infestation from infested trees to healthy stands which means the elimination of the trees that are already infested, in order to prevent cross-contamination. Ideally, this selective felling should be coupled with the thinning of the adjacent areas of healthy trees to help strengthen them and to prevent any wildfires from reaching the tree canopies through excessive ladder fuels. Again, this operation should be considered obligatory for wildland-urban interface zones, national and state parks and other areas frequented by the public for recreational purposes or travel.

This practical approach requires the collection and disposal of the felled trees in such a manner as to guarantee the total destruction of all living beetles and larvae within or attached to the trees and associated wood debris. Numerous methods have been employed towards this goal, including burning the trees in an above-ground air curtain burner (*FireBox*)⁸, chipping the trees

and hauling the residue to a landfill or biomass cogeneration facility, composting the trees after grinding them, and even salvaging portions for firewood or commercial uses.

Tree Disposal Considerations

As already stated, beetle infested trees must be slated for disposal for two main reasons:

1. To prevent spreading the infestation to healthy trees and
2. To prevent or mitigate forest fires, as the dead trees would be fuels for devastating wildfires that would likely spread to healthy stands with high ladder fuels.

Time is always of the essence regarding either objective. Beetle infested trees and slash must be effectively disposed of in the shortest amount of time to keep live beetles from migrating to healthy trees and larvae from developing into mature beetles which would then fly to healthy stands. This should best take place in winter and early spring before the larvae typically hatch.

Forest patches of dead trees could be ignited by lightning, fallen rocks or human activity at any time with possibly devastating consequences. The disposal of these trees could be performed year-round.

Above-ground air curtain burners, such as the portable models S-116, S-220 or S-327 fireboxes by Air Burners, Inc., have been designed for the disposal of clean wood waste (vegetative waste) as an alternative to open burning. Describing how an air curtain burner works is beyond the scope of this paper and the reader is directed to a technical memorandum on the principle of operation of air curtain incineration at this Web Site: <http://www.air-burnertechnology.com>.

Air curtain FireBoxes are the most desirable and suitable machines to accomplish the disposal of beetle infested trees for the following main reasons⁸:

1. The attained high burn temperatures assure quick and total elimination of any and all beetles and larvae in or on the felled tree and collected slash.
2. The wood debris can be burned immediately upon collection, even while the freshly cut tree is still *green*. A drying-out period is not required. That gives no opportunity to any larvae population in the tree to mature into beetles that would fly away and infest healthy trees.

3. Large sections of tree trunks and brush can be loaded without any or any excessive milling, avoiding the attraction of beetles from the release of conifer resins that may affect beetle behavior as the resins resemble beetle pheromones⁷.
4. The air curtain burner achieves 97-99% mass reduction and the resultant ash residue can almost always be applied to the land on site. This eliminates any hauling by trucks.
5. The air curtain burner provides the most cost-effective solution for the disposal of wood waste, both from the capital investment angle and the direct operating costs (see last section) and it has a useful life of 10-15 years.
6. The air curtain burner is environmentally friendly and its implementation has a limited operational "carbon footprint" in comparison with other disposal methods, as it only employs a small Diesel engine.
7. The air curtain FireBox meets or exceeds US EPA regulations for air curtain incinerators.
8. The air curtain burner is batch loaded, is simple to operate without a dedicated attendant and has virtually no downtime for repairs.
9. The air curtain burner is portable, delivered fully assembled and it can be relocated on site simply by dragging it on its skids.

All alternative disposal options have serious drawbacks.

Chipping was historically considered the preferred option and it was advocated that all beetles and larvae would be 100% destroyed by the violent process within the grinding and chipping machines powered by huge engines. It was thought also that chipping would be the most environmentally friendly alternative and the most economical, as biomass co-generation plants could turn the beetle infested trees into electric power. These premises turned out to be mostly false.

Small-scale tests were carried out by Deborah Mc McCullough, et al, of Michigan State University in 2003 to verify that all Emerald Ash Borer (EAB) beetles and larvae would be killed, if chipping resulted in chips smaller than 1 inch (25.4mm) long⁹. The typical size of this beetle is about one half inch (13.5mm) and its larvae slightly more than 1 inch (32mm)¹⁰ in length. The small-scale test did not appear to represent the real-life picture. Beetles and larvae do survive the chipping process. One such report in support of this finding, also from Michigan, shows that the Emerald Ash Borer infested stands of elm trees in circles around a biomass co-generation facility to which chips from Emerald Ash Borer infested trees were hauled by trucks¹¹. This was later verified by a 2005 study conducted by David L. Roberts, et al, of Michigan State University Extension¹².

Another mistake often made is to refer to the Michigan State University McCullough study in order to justify that chipping trees infested with the Mountain Pine Beetle (or similar) into chips of one inch in size will suffice to kill also this beetle and its larvae. What is not considered is the fact that the mountain pine beetle is much smaller than the Emerald Ash Borer that was used in the Michigan sample. The mountain pine beetle or its larvae is typically less than 1/5 inch (5mm) long¹³. Applying the assumptions of the McCullough study would require that trees infested with the mountain pine beetle be ground into chips smaller than about 1/10 inch (2.5mm), in order to kill the beetles and larvae effectively. Even then, it is very doubtful that all the tiny beetles and larvae would actually be hit by the cutting mechanism of the chipper or grinder. Chipping to such a small chip size would usually require more than one pass and is not practical and economical; and it is not what is actually being observed in the field today.

Also, chippers do not handle freshly cut "green" trees and brush very well, although that must be a requisite for effective beetle control. The wood waste drying time that would be required for effective chipping may give larvae ample time to mature and fly off to infest healthy trees. Another reason why the chipping and grinding of green trees would not be advisable is the fact that the chipping causes the release of large amounts of conifer resins in volatile form that attract beetles. This tends to lead to cross colonization, as the infested taken down green trees would usually be close to "leave" trees, the ones to be saved in selective felling initiatives.

Another problem plaguing the chipping operators is the fact that the chips cannot be indiscriminately applied to the forest floor on site. Chips on the forest floor are unnatural and adversely affect the forest ecosystem; that is why the layer of chips that is acceptable is limited by forest scientists. As a consequence the chipped trees will have to be hauled to a landfill at considerable cost and, again, possibly causing cross contamination on the way. Usually the chips cannot be sent to biomass cogeneration plants, because either the chip specifications are not acceptable or the transport costs to a suitable facility are too high.

Finally, a chipper is actually not as environmentally friendly as often proclaimed. The emissions from the massive chipper Diesel engine and the hauling trucks coupled with the (carcinogenic) wood dust released have a greater negative impact on the environment than air curtain burners. Air curtain burners use a small Diesel engine that is fuel efficient and the burning of clean wood is actually a natural process that has occurred on earth for millions of years. Also, the overall cost of the chipper operation is much higher when measured against air curtain burning, as will be demonstrated later.

The remaining tree disposal options include open pile burning, hauling to and depositing the wood debris into a landfill (usually after chipping the wood debris first to make hauling more manageable) or to compost it which also requires grinding. Little or no guarantee is provided, however, that cross-contamination is prevented during the transport over public roads and at the landfill or composting site. Composting itself has most of the drawback of chipping and more and would be the least attractive option. Open burning is usually not an option, as the burning of green trees which is difficult in itself would have to take place in close proximity to "leave" trees, as transporting the wood debris off-site would obviously be counterproductive. Smoke from open pile burning of wood waste is a serious issue. Entire valleys are known to be filled with smoke for days at a time. Open burning presents a serious wildfire danger, as a "controlled burn" often leads to an out-of-control forest fire. The May 2000 Cerro Grande Fire of Los Alamos, New Mexico, is a testimony of that. It started with a sanctioned prescribed burn and turned into a wildfire causing devastating losses. Interestingly enough, Los Alamos National Laboratory acquired several air burners, including the large S-327 FireBox. They were used for several years in post wildfire clean up and forest fire mitigation work after that disaster.

Dead trees from a beetle kill that comprise large areas should be removed for wildfire prevention and rehabilitation (reforestation) as they pose a serious fire danger. In this wood waste disposal application it is the cost effectiveness alone that makes air curtain burners more desirable than any other disposal option, except possibly, open pile-burning where permitted and if the smoke impact is not considered a nuisance or hazard as already addressed.

More About Air Curtain Burners¹⁴


Above-ground refractory walled [FireBoxes](#) by Air Burners, Inc. have a useful life of 10 to 15 years, making them a prudent capital acquisition. Depending on the model, direct operating costs are between \$7.00 and \$10.00 per hour based on late 2017 Diesel fuel cost, and the operation of the machines is simple and safe. Ancillary equipment for loading the FireBoxes can range from excavators to front loaders with a rake and even Bobcats, heavy equipment that usually is already on hand with experienced operators in the forest industry. Depending on the model chosen, Air Burners' FireBoxes can be transported on tilt-bed trailers, low-boys, drop decks, etc., or by the custom slide-axle trailers from *Landoll* that feature self-loading and unloading of all FireBoxes.

All Air Burners, Inc. air curtain burners are delivered completely assembled ready for immediate use. There is no need for set-up or tear down at all.

The fireboxes are designed to be dragged on their skids and they have no bottoms, as they are placed directly onto the ground.

The 2017 cost range of Air Burners' standard above-ground air curtain burners is roughly \$80,000 to \$165,000 (which includes the smaller trailer-mounted "[BurnBoss](#)", but not the electric power generation [PGFireBox](#)), and all machines are manufactured in the USA at the Air Burners, Inc. factory in Palm City, Florida. All units meet the US-EPA regulations for air curtain incinerators¹⁵.

New products include selected FireBoxes adapted so that they can be loaded and unloaded by standard roll-off trucks as are used throughout the US for many years ("[Roll-off FireBox](#)"). The principle of loading and unloading will mirror that of standard 20-foot construction dumpsters or cans. This will further simplify the deployment of fireboxes to the landings in the deep forest and the collection sites in the wildland-urban interface zones. A single FireBox could also be easily shared by several communal entities and transported by equipment that is usually already available commercially in the local area.



S-220 Firebox



S-220 Firebox

Summary

Many forests in North America are being destroyed by a variety of pests, mostly beetles. One major cause is related to the weakening of the trees due to the buildup of forest understory for nearly 100 years which has not been cleared by naturally occurring low-severity forest fires. These weakened stands are thus susceptible to beetle infestation. Trees killed by beetles must be removed to prevent the spreading of beetle infestations and as a wildfire mitigation and restoration effort. Beetle infested trees removed in selective felling operations must best be destroyed on site and while still green. The most effective and most commercially viable method for the disposal of these trees is the use of above-ground air curtain burners also called FireBoxes by

Air Burners, Inc. They are used, for example, by the US National Park Service, the USDA Forest Service, State of California and the Rocky Mountain national Park in Colorado and many other government agencies and private forest industry businesses in the US and Canada. The patented above-ground air curtain burners are manufactured by Air Burners, Inc. of Palm City, Florida and available from the factory.

References:

1. CBC News, "*Pine beetle outbreak adds to greenhouse gas woes*," April 2008, accessible at <http://www.cbc.ca/news/technology/pine-beetle-outbreak-adds-to-greenhouse-gas-woes-1.725982>
2. The Champion Tree Project, "*The Magic Tree*," USDA supported Educational Program, accessible at <http://www.youtube.com/watch?v=TcnWLXz0iVg>
3. Dean Lueck, Jonathan Yoder "*Clearing the Smoke From Wildfire Policy: An Economic Perspective* accessible at <http://www.perc.org/sites/default/files/pdfs/PS-56-WEB%20final.pdf>
4. "*CAL FIRE – Benefits of Fire*," accessible at http://www.fire.ca.gov/communications/downloads/fact_sheets/TheBenefitsofFire.pdf
5. "*Sequoia Sempervirens: Trial by Fire*," by Eileen Jones accessible at http://www.wildcarebayarea.org/site/PageServer?pagename=eNews_August2012_Redwoods
6. Colorado State University, "*Mountain Pine Beetle*," accessible at <http://extension.colostate.edu/topic-areas/insects/mountain-pine-beetle-5-528/>
7. Leatherman, Dave, "*Fire Fuels Reduction and Bark Beetles*," Colorado State Forest service, October 2002, accessible at http://www.frftp.org/sites/default/files/education/bark_beetles_fuel_mitigation.pdf
8. Air Burners, Inc., "*Above –ground Air Curtain Burners (FireBoxes)*," accessible at <https://airburners.com/products/firebox-series/>
9. US Forest Service Northern Research Station, "*Effects of chipping, grinding, and heat on survival of emerald ash borer, Agrilus planipennis (Coleoptera: Buprestidae)*," McCullough, Deborah G., et al, 2003, accessible at

<http://www.treesearch.fs.fed.us/pubs/12553> or
http://www.nrs.fs.fed.us/pubs/jrnl/2003/nc_2003_mccullough_002.pdf

10. United States Department of Agriculture Forest Service
Northeastern Area State and Private Forestry NA-PR-02-04, "*Pest Alert*", January 2004, accessible at
http://www.na.fs.fed.us/spfo/pubs/pest_al/eab/eab04.htm

11. The Detroit News" Special Report: "*Flaws riddle ash borer fight, Quarantine, grinding fail to contain beetles in Michigan,*" October 2004, no longer accessible at <http://www.wzzm13.com/news/article/32069/0/Flaws-riddle-ash-borer-fight>

12. Michigan State University Extension: "*The Survival of EAB in Wood Chips: Does Size Matter?*", Roberts, David L., et al, 2005, no longer accessible at http://treedoctor.anr.msu.edu/ash/Roberts_ash%20woodchips.pdf

13. Wikipedia, "*Mountain Pine Beetle*", accessible at https://en.wikipedia.org/wiki/Mountain_pine_beetle

14. Pearl harbor – Hickam News, "*Base brings burner to beetle battle to help prevent spread,*" October 2014 accessible at
<http://www.hookelenews.com/base-brings-burner-to-beetle-battle-to-help-prevent-spread/>

15. US EPA Debris reduction of vegetative material video, July 2008, accessible at <https://airburners.com/resource-center/videos/>

Norbert Fuhrmann is a free-lance writer and researcher. He holds an M.B.A. from the Crummer Graduate School of Business at Rollins College in Winter Park, Florida.

* * * * *